



**QUEEN'S
UNIVERSITY
BELFAST**

Lipopolysaccharide modification in Gram-negative bacteria during chronic infection

Maldonado, R. F., Sá-Correia, I., & Valvano, M. A. (2016). Lipopolysaccharide modification in Gram-negative bacteria during chronic infection. *FEMS Microbiol. Rev.*, 40(4), 480-493. <https://doi.org/10.1093/femsre/fuw007>

Published in:
FEMS Microbiol. Rev.

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

©2016 Oxford University Press.

This is a pre-copyedited, author-produced PDF of an article accepted for publication in FEMS Microbiology Reviews following peer review. The version of record is available online at: <http://femsre.oxfordjournals.org/content/40/4/480>

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Lipopolysaccharide modification in Gram-negative bacteria during chronic infection

Rita F. Maldonado¹, Isabel Sá-Correia¹, and Miguel A. Valvano^{2,3,*}

¹Institute for Bioengineering and Biosciences, Department of Bioengineering, Instituto Superior Técnico, Universidade de Lisboa, Lisbon, 1049-001, Portugal, ²Department of Microbiology and Immunology, University of Western Ontario, London, Ontario, N6A 5C1, Canada and ³Centre for Infection and Immunity, Queen's University Belfast, Belfast, BT9 7BL, United Kingdom.

***Corresponding author:** Centre for Infection and Immunity, Queen's University Belfast, 97
Lisburn Rd., Belfast, BT9 7BL, United Kingdom. Tel: +44-28-9097-6025; E-mail:
m.valvano@qub.ac.uk

One sentence summary: We review modifications of lipopolysaccharide structure and biosynthetic pathways that occur upon bacterial adaptation to chronic respiratory and gastrointestinal infections.

Keywords (6): adaptive mutation, O antigen, lipid A, *Pseudomonas aeruginosa*, *Burkholderia cenocepacia*, cystic fibrosis, *Helicobacter pylori*, gastric ulcer

Running title: Lipopolysaccharide modification during chronic infection

ABSTRACT

The Gram-negative bacterial lipopolysaccharide (LPS) is a major component of the outer membrane that plays a key role in host-pathogen interactions with the innate immune system. During infection, bacteria are exposed to a host environment that is typically dominated by inflammatory cells and soluble factors, including antibiotics, which provide cues about regulation of gene expression. Bacterial adaptive changes including modulation of LPS synthesis and structure are a conserved theme in infections, irrespective of the type or bacteria or the site of infection. In general, these changes result in immune system evasion, persisting inflammation, and increased antimicrobial resistance. Here, we review the modifications of LPS structure and biosynthetic pathways that occur upon adaptation of model opportunistic pathogens (*Pseudomonas aeruginosa*, *Burkholderia cepacia* complex bacteria, *Helicobacter pylori* and *Salmonella enterica*) to chronic infection in respiratory and gastrointestinal sites. We also discuss the molecular mechanisms of these variations and their role in the host-pathogen interaction.

INTRODUCTION

The lipopolysaccharide (LPS) is a central component of the outer membrane in Gram-negative bacteria and frequently plays a key role in pathogenesis (Figure 1) (Whitfield & Trent, 2014). LPS is the dominant glycolipid in the outer leaflet of the outer membrane, forming a layer that is stabilized by divalent cations and provides an effective permeability barrier against deleterious molecules such as antibiotics and cationic antimicrobial peptides (Nikaido, 2003). The classical LPS molecule has a tripartite structure comprising (i) lipid A, the hydrophobic moiety that anchors LPS to the outer leaflet of the outer membrane, (ii) core oligosaccharide (herein core), which together with lipid A, contributes to maintain the integrity of the outer membrane, and (iii) O antigen polysaccharide or O antigen, which is connected to the core and consists of a polymer made of repeating oligosaccharide units in direct contact with the external milieu (Figure 1) (Whitfield & Trent, 2014). LPS molecules only including lipid A and core are generally referred to as "rough" and often called lipooligosaccharides, while the complete LPS capped with O antigen is called "smooth".

The lipid A is embedded in the outer membrane and composed of acyl chains linked to

a backbone dimer of glucosamine by ester and/or amide linkages. The typically hexa-acylated lipid A elicits robust inflammatory responses upon recognition by the complex Toll-like receptor 4 and myeloid differentiation factor 2 (TLR4-MD2), which is predominantly found on macrophages, monocytes and dendritic cells (Park *et al.*, 2009, Park & Lee, 2013).

Modification of the lipid A acylation patterns, or addition of positively-charged substituents to the lipid A phosphate groups (Raetz *et al.*, 2007), confer protection against host innate defenses by reducing even further the permeability of the outer membrane to antimicrobial peptides and dampening inflammatory responses by the host (Raetz *et al.*, 2007, Needham & Trent, 2013, Di Lorenzo *et al.*, 2015).

Lipid A is glycosylated at the 6'-position with two residues of 3-deoxy-D-manno-oct-2-ulosonic acid (Kdo); the inner Kdo serves as the point of attachment for the remaining core. Some bacterial species such as *Burkholderia* (Silipo *et al.*, 2005, Silipo *et al.*, 2007) produce a modified Kdo, which is converted into D-glycero-D-talo-oct-2-ulosonic acid (Ko) by a unique Kdo-3 hydroxylase (Chung & Raetz, 2011). The next sugars added to the Lipid A-Kdo₂ are typically two or more residues of L-glycero-D-manno-heptose, although in some species LPS molecules are devoid of heptose (Valvano *et al.*, 2002). The rest of the core consists of a set of sugars that differs among species and even among strains of the same species (Mamat *et al.*, 2011). Phosphorylation of the core sugars in *P. aeruginosa* has been associated with increased membrane impermeability and resistance to antibiotics (Walsh *et al.*, 2000), and is also required for the transport of LPS to the outer membrane (Delucia *et al.*, 2011). The *P. aeruginosa* core may also be a ligand for the cystic fibrosis (CF) transmembrane conductance regulator protein displayed on the apical surface of epithelial cells (Schroeder *et al.*, 2002).

O antigens comprise repeating oligosaccharide units that may be linear or branched (Whitfield & Trent, 2014). The O-repeating unit is highly variable immunochemically giving rise to a vast number of different O-specific serotypes (Valvano *et al.*, 2011, Whitfield & Trent, 2014). The O antigen contributes to evasion of host immune defenses, particularly evasion of the complement cascade in *Salmonella enterica* serovar Typhimurium (Murray *et al.*, 2006), delay of recognition and internalization in epithelial cells in *Salmonella* Typhimurium and *Burkholderia cenocepacia* (Duerr *et al.*, 2009, Saldías *et al.*, 2009), enhanced intracellular survival in *Shigella flexneri* (West *et al.*, 2005) and *Brucella melitensis* (Paixão *et al.*, 2009),

86 and protection against oxidative stress in *Erwinia amylovora* (Berry *et al.*, 2009). O antigen
87 also contributes to swimming and swarming motility in *E. amylovora* (Berry *et al.*, 2009), *B.*
88 *cenoeopacia* (Coutinho *et al.*, 2011), and *Pectobacterium atrosepticum* (Bowden *et al.*, 2013).
89 The immunogenicity of the O antigen polysaccharide elicits a robust antibody response, which
90 may cause selective pressure on bacteria to lose the ability to produce O antigen (King *et al.*,
91 2009); this is particularly common for chronic *P. aeruginosa* strains infecting the lungs of CF
92 patients (Hancock *et al.*, 1983). Conceivably, once the bacteria become mucoid (Govan &
93 Deretic, 1996), the nutrient burden is so high producing alginate and that the bacteria are
94 replicating in a “protected” niche in which O antigen becomes dispensable. However, this may
95 not be a universal notion since other bacteria chronically infecting the CF lung, such as
96 members of the *B. cepacia* complex, undergo different adaptive changes than those reported
97 for *P. aeruginosa* (Zlosnik *et al.*, 2014), including the observation of an inverse correlation
98 between the quantity of mucoid exopolysaccharide production and the rate of decline in CF
99 lung function (Zlosnik *et al.*, 2011).

100 Most *P. aeruginosa* strains produce two types of O antigen molecules (“A-band” and
101 “B-band”), which are structurally and serologically distinct and have different mechanisms of
102 biosynthesis (King *et al.*, 2009, Lam *et al.*, 2011). The “A-band” or “common polysaccharide
103 antigen” is a homopolymer of D-rhamnose that elicits a relatively weak antibody response
104 (King *et al.*, 2009). “B-band” or “O-specific antigens” are highly immunogenic heteropolymers
105 composed repetitive units of different sugars and form the basis for the AITS *P. aeruginosa*-
106 serotyping scheme that includes 20 serotypes (Knirel *et al.*, 2006). Structural data in several
107 *Pseudomonas* serotype strains (Sadovskaya *et al.*, 2000, Bystrova *et al.*, 2006) and genetic
108 experiments (Abeyrathne *et al.*, 2005) demonstrate that both common and O-specific
109 antigens are linked to the lipid A-core.

110 In this article, we review the literature on LPS variations occurring upon bacterial
111 adaptation to chronic infection, with special emphasis on chronic respiratory infections in
112 patients with CF and gastric infections. CF is a genetic disease that leads to ineffective
113 mucociliary clearance of the airways, resulting in chronic airways infection by several Gram-
114 negative bacterial opportunistic pathogens, such as *P. aeruginosa*, the *Burkholderia cepacia*
115 complex (Bcc), and *Achromobacter xylosoxidans* (Ciofu *et al.*, 2015, Cullen & McClean, 2015,

Parkins & Floto, 2015). Chronic gastric infection by *H. pylori* leads to a pre-cancerous state associated with loss of acid-producing parietal cells, which results in increased gastric pH, and pepsinogen-producing zymogenic cells. The gastric environment changes during disease progression and as a result, infecting *H. pylori* strains must adapt to persist in a gastric habitat with increased pH and different cell composition (Skoglund *et al.*, 2009, Rubin & Trent, 2013, Malnick *et al.*, 2014). Because both respiratory infections in CF patients and gastric infections by *H. pylori* remain during the lifetime of the patient, they provide natural human models of disease progression and microbial adaptation to the host environment.

LPS BIOSYNTHESIS

Lipid A-core biosynthesis

The biosynthesis of LPS has been reviewed in detail elsewhere (Raetz *et al.*, 2007) (King *et al.*, 2009, Lam *et al.*, 2011, Greenfield & Whitfield, 2012, Whitfield & Trent, 2014, Valvano, 2015). Briefly, the lipid A is synthesized on the cytoplasmic side of the inner membrane by a conserved pathway of nine enzymes catalyzing the sequential conversion of the precursor UDP-*N*-acetyl-glucosamine into lipid A-Kdo₂, which is the acceptor for the rest of the core sugars that are added from nucleotide sugar precursors *via* sequential glycosyl transfer reactions (Figure 2) (Mamat *et al.*, 2011, Whitfield & Trent, 2014). The complete lipid A-core is transported to the periplasmic face of the inner membrane by the ABC transporter MsbA (Whitfield & Trent, 2014). Diverse covalent modifications of lipid A may occur during its transit from the periplasmic side of the inner membrane to the outer leaflet of the outer membrane (Raetz *et al.*, 2007), which are important for niche adaptation and can influence the virulence of the pathogen (Needham & Trent, 2013). In bacteria that produce O antigen, the O polysaccharide is assembled by a separate biosynthesis pathway (see next section) and attached to the core at the periplasmic side of the inner membrane (Figure 2).

O antigen biosynthesis

The O antigen is synthesized by cytoplasmic membrane-associated enzyme complexes and requires C₅₅-undecaprenyl phosphate (Und-P), which serves as an acceptor for O antigen

chain assembly (Valvano, 2011). Chain assembly occurs by the action of diverse glycosyltransferases that synthesize the specific O antigen of each strain. Genes at the *wb** (formerly *rfb*) locus encode most of the enzymes involved in O antigen assembly (Reeves *et al.*, 1996). Because of the great diversity of O antigen structures, the *wb** loci are highly polymorphic (Raetz & Whitfield, 2002, Lam *et al.*, 2011). The O antigen is initially assembled on the cytoplasmic side of the membrane and then translocated to the periplasmic side and ligated to lipid A-core (Valvano, 2015) (Figure 2). There are three pathways for O antigen biosynthesis and export: (1) Wzy-dependent, (2) ABC-transporter-dependent, and (3) synthase-dependent (Keenleyside & Whitfield, 1996, Lam *et al.*, 2011, Greenfield & Whitfield, 2012, Valvano, 2015). The mature LPS molecule is then transported across the periplasm and inserted into the outer leaflet of the outer membrane by the conserved Lpt (LPS transport) pathway (May *et al.*, 2015, Simpson *et al.*, 2015). Lpt proteins form a complex that traverses the Gram-negative cell envelope to deliver LPS to the outer membrane and include an ABC protein complex (LptBFG) that uses energy from ATP hydrolysis to extract LPS from the periplasmic face of the inner membrane, several proteins that dock and promote the transfer of LPS across the periplasm (LptCA and YhjD) and a complex of proteins on the outer membrane (LptDE, YtfN, YfgH and YceK), responsible for the correct insertion of LPS in the outer leaflet (Babu *et al.*, 2011, Sperandio *et al.*, 2011, Sperandio *et al.*, 2011, May *et al.*, 2015, Simpson *et al.*, 2015). The Lpt system has not been investigated in Gram-negative pathogens other than *E. coli* and sequence homology between *E. coli* and *P. aeruginosa* genes is low, with the exception of LptB (66% sequence identity). Recently, it was shown that *P. aeruginosa* LptA has a dimeric structure, unlike the oligomeric structure of *E. coli* LptA (Shapiro *et al.*, 2014).

In *P. aeruginosa*, the common polysaccharide and the O-specific antigens are synthesized *via* the ABC-transporter-dependent pathway and the Wzy-dependent pathway, respectively (King *et al.*, 2009, Lam *et al.*, 2011). In both the synthesis is initiated by the same glycosyltransferase, WbpL (homologous to the *E. coli* WecA), resulting the formation of an Und-P-P-sugar intermediate (King *et al.*, 2009, Lam *et al.*, 2011). Four enzymes are required for the biosynthesis of GDP-D-rhamnose, the nucleotide sugar precursor for the common polysaccharide antigen: WbpW, AlgC, Gmd and Rmd (King *et al.*, 2009, Lam *et al.*, 2011).

The glycosyltransferases WbpX, WbpY and WbpZ are involved in the synthesis of the common polysaccharide antigen (King *et al.*, 2009, Lam *et al.*, 2011), while genes *pa54-55pa5459* have been suggested to encode proteins that play a role in controlling chain length (Hao *et al.*, 2013). Once the common polysaccharide antigen is linked to the Und-P carrier, the complex is exported across the membrane by the ABC-transport system Wzm-Wzt (King *et al.*, 2009, Lam *et al.*, 2011). While the genes for the synthesis and assembly of the common polysaccharide are conserved, different set of genes are responsible for the biosynthesis of the O-specific antigen in each serotype strain. These genes are in a cluster flanked by the highly conserved genes *himD/ihfB* and *wbpM* (King *et al.*, 2009, Lam *et al.*, 2011). While the *P. aeruginosa* O5, O6 and O11 O antigen clusters were studied to some extent very little experimental work was conducted into the functions of genes in the remaining O antigen loci (Lam *et al.*, 2011). The synthesized the Und-PP-linked O-repeat units are translocated to the periplasmic side of the membrane and polymerized. The proteins Wzy, Wzz and Wzx are required for this process, acting as polymerase, chain-length regulator, and flippase, respectively (Lam *et al.*, 2011). Once on the periplasmic side, both the common polysaccharide antigen and the O-specific antigen are independently linked to the lipid A-core complex by the WaaL ligase (Figure 2) (Abeyrathne *et al.*, 2005, Valvano, 2011, Ruan *et al.*, 2012).

LPS VARIATION DURING CHRONIC RESPIRATORY INFECTIONS IN CF PATIENTS

***Pseudomonas aeruginosa* infection**

P. aeruginosa is the most common pathogen isolated from the respiratory tract of adult patients with CF (Lipuma, 2010, Hauser *et al.*, 2011). Chronic airway infections caused by *P. aeruginosa* are found in up to 80% of adult patients with CF (Aaron *et al.*, 2010, Lipuma, 2010) and are associated with increased morbidity and mortality (Hauser *et al.*, 2011). Phenotypic changes suggesting *P. aeruginosa* adaptation to the CF lung have been reported in several studies (Hogardt & Heesemann, 2010). They include loss of motility associated with growth in microcolony (Sriramulu *et al.*, 2005), reduced expression of virulence factors, which is presumably an adaptive strategy to escape detection by the host immune system (Smith *et*

al., 2006), increased activity of efflux pumps associated with antibiotic resistance, especially against those antibiotics used clinically (Poole, 2005), and a switch from non mucoid to mucoid phenotypes (Ciofu *et al.*, 2010, Hogardt & Heesemann, 2010). The phenotypic changes reflect point mutations accumulating in *P. aeruginosa* lineages that persist in CF airways (Lorè *et al.*, 2012), and include mutations in alginate biosynthesis regulator genes (Bragonzi *et al.*, 2006) and genes involved in the LPS modification (Cigana *et al.*, 2009), motility (Mahenthiralingam *et al.*, 1994), quorum-sensing regulation (D'Argenio *et al.*, 2007, Hoffman *et al.*, 2009), type 3 secretion system biosynthesis (Jain *et al.*, 2004), multidrug-efflux pumps, and mutator genes (Oliver *et al.*, 2000).

The longitudinal course of chronic airway infection with *P. aeruginosa* in CF has been followed in various studies (Smith *et al.*, 2006, Cigana *et al.*, 2009, Cramer *et al.*, 2011, Mowat *et al.*, 2011, Warren *et al.*, 2011, Yang *et al.*, 2011, Lorè *et al.*, 2012, Dettman *et al.*, 2013). A study investigating over 1700 serial isolates obtained from 10 patients infected with the same strain showed that within-patient diversity made the largest contribution to the overall variation in the population and also that population compositions fluctuated over time (Mowat *et al.*, 2011). The authors suggested that extensive diversity within the *P. aeruginosa* population during chronic infection has the potential to provide a reservoir for antibiotic resistant mutations and mutations in other virulence traits (Mowat *et al.*, 2011). Despite these differences, certain traits were overrepresented in all isolates, most of which include properties regulated by quorum sensing (Mowat *et al.*, 2011). *In silico* simulations reveal virulence factor expression decline towards the end of chronic infections and adaptive mutations that tend to improve metabolic fitness, which would optimize growth over the more energetically expensive virulence factor production (Oberhardt *et al.*, 2010). *P. aeruginosa* LPS modifications appear to be an important factor in the adaptation of this pathogen to chronic infection (Cigana *et al.*, 2009). Indeed, chronic *P. aeruginosa* CF isolates have rough colony phenotypes and contain few, short, or no O side chains, becoming non-typeable (Hancock *et al.*, 1983). O antigen deficient isolates are sensitive to *in vitro* killing by serum complement and become more tolerant to the antibiotic gentamicin (Kadurugamuwa *et al.*, 1993). Analysis of sequential variants of *P. aeruginosa* show O antigen loss (Lee *et al.*, 2005) and lipid A modifications (Cigana *et al.*, 2009). Whole-genome analysis of two clinical *P.*

aeruginosa variants recovered from a chronic CF patient after 6 and 96 months of infection also revealed non-synonymous mutations in the O antigen biosynthetic genes *wbpA* and *pa5238* in the latter variant (Smith *et al.*, 2006). Another study of genes responsible for modifying lipid A revealed one mutation in *pagL* in late variants, which abolish PagL expression and leads to reduced TLR4-MD2-signalling (Cigana *et al.*, 2009). Thus, initial lipid A modifications by addition of palmitate to the lipid A of *P. aeruginosa* make the LPS more proinflammatory, but the subsequent modification through the loss of PagL activity decreases its proinflammatory activity. Together, the results of these studies suggest that reduced LPS immunostimulatory potential contributes to immune system evasion and survival over the course of the chronic *P. aeruginosa* infection. Experimental data support this hypothesis since a comparison of the pathogenicity of nine *P. aeruginosa* sequential clonal variants in the infection models *Caenorhabditis elegans*, *Galleria mellonella*, *Drosophila melanogaster* and two different mice backgrounds (C57Bl/6NCrI and BALB/cAnNCrI), show that early *P. aeruginosa* variants were lethal in all infection models tested, while late strains exhibited reduced or no virulence (Lorè *et al.*, 2012).

A microevolution analysis based on whole-genome sequencing of sequential *P. aeruginosa* variants recovered from CF patients for more than 20 years (Cramer *et al.*, 2011) identified codon changes in genes for lipid A biosynthesis (*lpxC*, *lpxO2* and *yciK*), core biosynthesis (*rfaD* and *wapP*), and common polysaccharide antigen biosynthesis (*wbpZ*) (Cramer *et al.*, 2011). Another genomic analysis taken over 200,000 bacterial generations of 12 selected *P. aeruginosa* DK2 variants recovered from six CF patients identified a total of 234 non-synonymous single nucleotide polymorphisms among the genomes in relation to their common ancestor strain, suggesting that an initial period of rapid adaptation is followed by a period of genetic drift in this lineage (Yang *et al.*, 2011). Three of the non-synonymous single nucleotide polymorphisms occurred in genes needed for lipid A biosynthesis and modification (*pagL* and *lpxO2*) and O-specific antigen synthesis (*wzz*) (Yang *et al.*, 2011). A recent study analyzing whole-genome sequence data from *P. aeruginosa* clinical isolates sampled from the sputum of 32 different patients reported that the O antigen ligase *waaL* is one of the few hotspots of gene polymorphisms (Dettman *et al.*, 2013). To gain insight into the role of mutator genes for generating adaptive variation, Warren *et al.* (2011) analysed the

genomes of two series of isolates recovered from two patients, similar in duration but different in mutator incidence, and identified 15 LPS genes that lacked in multiple members both in mutator and non-mutator series. All the identified genes are involved in the synthesis of serogroup O2/O5/O26/O18/O20 O antigen (*wbpA*, *wbpB*, *wbpC*, *wbpD*, *wbpE*, *wbpG*, *wbpH*, *wbpI*, *wbpJ*, *wbpK*, *wbpL*, *wzx*, *wzy*, *wzz* and *pa1385*) (Warren *et al.*, 2011).

In addition to changes in O antigen, adaption of *P. aeruginosa* to chronic lung infection in CF patients involves the synthesis of various lipid A structures (Figure 3) (Ernst *et al.*, 2007), which result in alteration of host innate immune responses and promote bacterial persistence (Moskowitz & Ernst, 2010). These modifications involve deacylation of the lipid A resulting in the loss of an acyl chain from the 3-position, which is catalyzed by PagL (Figure 3) (Trent *et al.*, 2001, Geurtsen *et al.*, 2005, Ernst *et al.*, 2006). Under acylation of lipid A has been associated with low inflammatory activity (Moskowitz & Ernst, 2010, Di Lorenzo *et al.*, 2015) and modulation of TLR4-MD2 receptor recognition (Ernst *et al.*, 2003). Also, *P. aeruginosa* lipid A can acquire a secondary acyl chain into the 3'-position, which is catalyzed by a divergent palmitoyltransferase functionally analogous to the *Salmonella* and *E. coli* PagP enzyme (Figure 3) (Thaipisuttikul *et al.*, 2014). Further modifications involve the addition of secondary acyl chains to the chains present at the 2- and 2'-positions, which is catalyzed by HtrB and LpxO, respectively (Figure 3), as well as the incorporation of 4-amino-4-deoxy-L-arabinopyranose (Arap4N) to phosphate groups at the 1- and 4'-positions by the two-component regulatory system PmrAB (Figure 3) (Moskowitz *et al.*, 2004). These lipid A modifications contribute to *P. aeruginosa* adaptation to the CF airway (Moskowitz & Ernst, 2010). The addition of phosphoethanolamine to the *P. aeruginosa* lipid A via the ColRS two-component system (Figure 3) in a Zn^{2+} -dependent manner was recently reported (Nowicki *et al.*, 2015), but the role of this modification *in vivo* is not clear.

Collectively, the studies described above support the notion that chronically infecting bacteria adapt to host immune responses by producing LPS lacking O antigen and by introducing lipid A modifications in isolates recovered in late stages of CF chronic infection (Table 1) (Lyczak *et al.*, 2002, Lee *et al.*, 2005, Smith *et al.*, 2006, Cigana *et al.*, 2009, Moskowitz & Ernst, 2010, Cramer *et al.*, 2011, Yang *et al.*, 2011, Dettman *et al.*, 2013). This conclusion is also supported from comparative studies using various host models

demonstrating that adaptation of different *P. aeruginosa* lineages within CF lungs selects populations with reduced pathogenic potential in acute infections (Lorè *et al.*, 2012).

Chronic Infections by other Gram-negative CF pathogens

Bacteria from the *Burkholderia cepacia* complex (Bcc) emerged as significant CF pathogens in the early 1980s, when a minority of infected patients exhibited rapid clinical deterioration, resulting in early death (Mahenthiralingam *et al.*, 2005, Loutet & Valvano, 2010). Respiratory infections with Bcc bacteria in CF patients generally lead to faster decline in lung function and, in some cases to cepacia syndrome, a fatal necrotizing pneumonia frequently accompanied by septicemia (Mahenthiralingam *et al.*, 2005, Coutinho *et al.*, 2011). Further, Bcc bacteria are transmissible through social contacts and are intrinsically resistant to most clinically used antibiotics, which renders their eradication from the CF lung very difficult, if not virtually impossible (Mahenthiralingam *et al.*, 2005, Drevinek & Mahenthiralingam, 2010, Coutinho *et al.*, 2011). Although transient infection of the respiratory tract may occur in some patients, acquisition of Bcc most typically results in chronic infection (Mahenthiralingam *et al.*, 2005, Coutinho *et al.*, 2011). The same level of adaptation is not so clear cut in *B. cenocepacia* infections, as studies using the various infection models (*C. elegans*, *G. mellonella*, alfalfa, mice and rats) reported that most virulence factors are specific for one infection model only and rarely essential for pathogenicity in multiple hosts (Uehlinger *et al.*, 2009, Lorè *et al.*, 2012). Furthermore, less is known about *Burkholderia* adaptation during CF chronic infection; however, there has been an effort to characterize the evolution of *Burkholderia* populations in the lung, including phenotyping (Coutinho *et al.*, 2011, Moreira *et al.*, 2014) and genotyping of serial isolates (Lieberman *et al.*, 2011, Traverse *et al.*, 2013), and comparative expression profiling of the transcriptome (Mira *et al.*, 2011) and the proteome (Madeira *et al.*, 2011, Madeira *et al.*, 2013).

Lieberman *et al.* (2011) sequenced the genomes of 112 clinical *Burkholderia dolosa* isolates that resulted from the evolution of a single strain in 14 CF patients over 16 years of epidemic spread and discovered that genes involved in oxygen regulation, antibiotic resistance, outer-membrane synthesis and secretion have recurrent mutation patterns (Lieberman *et al.*, 2011). Interestingly, recurrent mutations in the same amino acid of the

glycosyltransferase WbaD were observed in nine patients, which resulted in production of O-unit repeats that were absent in the ancestral phenotype (Lieberman *et al.*, 2011). The ancestral *B. dolosa* genotype encodes a stop codon at this locus that prevents O antigen synthesis. In some variants, two different mutations affecting the same amino acid were detected, both of them restoring the full-length WbaD protein and leading to O antigen production (Table 2) (Lieberman *et al.*, 2011). Although this gain-of-function mutation does not follow the loss of O antigen tendency described in *P. aeruginosa*, these results underpin the importance of the O antigen switch mechanism during chronic infection. Another metagenomic analysis of six lineages evolved in biofilm mode of growth revealed an extraordinary mutational parallelism, including genes known to affect LPS biosynthesis, transcription, galactose metabolism, tricarboxylic acid cycle enzymes and altered metabolism of cyclic diguanosine monophosphate (Traverse *et al.*, 2013). One commonly mutated locus, showing twenty independent mutations in both *B. dolosa* and *Burkholderia cenocepacia*, was *manC*, encoding a nucleotide mannose biosynthesis protein presumably involved in surface polysaccharide biosynthesis that could be either an exopolysaccharide or LPS (Traverse *et al.*, 2013). Interestingly, these authors showed that complementation of one of the *manC* mutations dramatically reduced biofilm formation, and they speculated that the loss of polysaccharide may be required for efficient biofilm formation rather than immune evasion (Traverse *et al.*, 2013).

A comparison of the transcriptome and the proteome of three *B. cenocepacia* isolates recovered at the beginning of the infection and later during the progress of the disease suggests that the expression from genes involved in LPS biosynthesis is altered during chronic infection (Madeira *et al.*, 2011, Mira *et al.*, 2011, Madeira *et al.*, 2013), in particular of those required for O antigen biosynthesis. Indeed, recent analysis of the LPS structure of these isolates revealed that, although the early-stage isolate has a complete LPS with the O-chain moiety, the late-stage variants have a rough-type LPS, lacking O antigen (Maldonado *et al.*, unpublished data).

Several studies at genome, transcriptome and proteome levels have contributed to a better understanding of Bcc bacteria genome-wide adaptive mechanisms during chronic infections. Together, they suggest that there is a high selective pressure on the O antigen

locus leading to alterations both at the structural, sequence and regulatory levels. Given the exceptional parallelism found among the relatively few studies dedicated to Bcc bacteria and *P. aeruginosa*, the LPS seems to play an important role during chronic infection, both in immune system evasion and biofilm adaptation. Moreover, lack of O antigen in *B. cenocepacia* leads to increased internalization into macrophages upon phagocytosis (Saldías *et al.*, 2009), which may explain the higher invasiveness of epidemic strains, such as J2315, which do not produce O antigen. O antigen loss could therefore facilitate access of Bcc bacteria to macrophages, where intracellular bacteria could find a niche to persist, in agreement with a recent study showing that in human lungs, Bcc bacteria but not *P. aeruginosa* are found mainly inside macrophages (Schwab *et al.*, 2014). Other Gram-negative opportunistic pathogens that cause CF chronic infections include *Stenotrophomonas maltophilia*, *Achromobacter xylosoxidans* and *Haemophilus influenza*. Recently, some studies characterizing the adaptive traits of sequential isolates of *S. maltophilia* (Vidigal *et al.*, 2014), *A. xylosoxidans* (Trancassini *et al.*, 2014) and *H. influenza* (Watson *et al.*, 2004) recovered from CF patients have been published, however the LPS characterization of these clinical isolates is still lacking.

LPS VARIATION DURING CHRONIC GASTRIC INFECTION

The human gastric pathogen *Helicobacter pylori* is usually acquired during childhood by colonizing the human gastric mucosa and producing a superficial gastritis, which may remain asymptomatic during the lifetime of colonized individuals or eventually lead to gastric ulcer and atrophic gastritis (Linz *et al.*, 2013, Otero *et al.*, 2014). This geographically wide-spread bacteria infects more than half of the human population and is one of the most genetically diverse bacterial species, being also one of the most ubiquitous infectious organisms (Linz *et al.*, 2013). The genetic diversity of *H. pylori* is caused by a high mutation rate, presumably due to the lack of several mutation repair genes (Kang & Blaser, 2006). Chronic infection with *H. pylori* is recognized as the most common cause of gastric and duodenal ulcers (Brown, 2000). *H. pylori* chronic infection is also associated with the development of gastric adenocarcinoma and lymphoma of mucosa-associated lymphoid tissue (Otero *et al.*, 2014,

385 Mégraud *et al.*, 2015), for which this bacterium is considered to be a class 1 carcinogen
386 (WHO, 1994).

387 *H. pylori* produces several virulence factors of which the vacuolating toxin A (VacA),
388 the cytotoxin-associated gene A (CagA), and LPS play major roles in immunomodulation and
389 contribute to maintain chronic infection (Posselt *et al.*, 2013, Rubin & Trent, 2013, Chmiela *et*
390 *al.*, 2014, de Bernard & Josenhans, 2014, Hatakeyama, 2014). These factors contribute to
391 maintain the infection by preventing the clearance of *H. pylori* from the gastric mucosa and
392 interfering with innate and adaptive immune responses. Structural modifications of the lipid A
393 result in reduced endotoxicity, while expression and variation of Lewis determinants exposed
394 on the bacterial cell surface as a terminal O-specific oligosaccharide (Aspinall *et al.*, 1996,
395 Monteiro *et al.*, 1998) mimic host components expressed on the human gastric epithelium
396 (Moran *et al.*, 1996, Moran, 2008) and reduce detection by the immune system. *H. pylori* lipid
397 A presents a unique structure and shows remarkably lower biological activity compared with
398 lipid A from other bacteria (Muotiala *et al.*, 1992, Moran & Aspinall, 1998). Structural analysis
399 revealed that the lipid A acyl chains are longer (16 to 18 carbons) than those present in
400 enterobacterial lipid A (Moran *et al.*, 1997). The predominant form is tetra-acylated lipid A,
401 which is also underphosphorylated (Moran *et al.*, 1997, Cullen *et al.*, 2011).

402 Underphosphorylation and underacylation of *H. pylori* lipid A are responsible for reduced
403 endotoxicity (Ljungh *et al.*, 1996), as determined by its low reactivity against anti-lipid A
404 antibodies (Mattsby-Baltzer *et al.*, 1992), reduced ability to induce the production of cytokines,
405 nitric oxide and prostaglandin E2 (Pérez-Pérez *et al.*, 1995), and E-selectin expression
406 (Darveau *et al.*, 1995), as well as reduced activation of leukocytes (Baker *et al.*, 1994,
407 Semeraro *et al.*, 1996). Lipid A remodelling in *H. pylori* occurs mainly on the periplasmic side
408 of the inner membrane. A first set of modifications involves removal of the 1-phosphate group
409 by LpxE and the addition of a phosphorylethanolamine in its place by EptA (Tran *et al.*, 2004,
410 Tran *et al.*, 2006). These modifications increase bacterial resistance to antimicrobial peptides
411 (Tran *et al.*, 2006). Second, a two-protein Kdo-hydrolase complex removes the terminal Kdo
412 sugar, a modification that is critical to allow the ligation of the O-specific oligosaccharides to
413 the lipid A core (Stead *et al.*, 2010). Third, LpxF catalyses the removal of the 4'-phosphate
414 group (Cullen *et al.*, 2011). After ligation of the O-specific oligosaccharide (see below) the

complete LPS molecule is transported and displayed on the surface of the bacterial outer membrane. Once in the outer membrane the lipid A undergoes a final modification that consist on the removal of the 3'-linked acyl chains by LpxR, producing the characteristic tetra-acylated lipid A structure (Stead *et al.*, 2008).

The *H. pylori* O-specific oligosaccharide is initially formed as a lipid-linked oligosaccharide resulting from the addition of monosaccharides, but does not form a repeating oligosaccharide unit (Berg *et al.*, 1997, Rubin & Trent, 2013). The O-specific oligosaccharide has a common backbone that is further modified by fucosyltransferases generating structures that mimic human Lewis antigen molecules and other related blood-group antigens such as Le^X, Le^Y, Le^a, Le^b, sialyl-Le^X, H-1 antigen, and blood groups A and B antigens (Rubin & Trent, 2013) (Figure 4) This lipid-linked fucosylated oligosaccharide is translocated across the inner membrane by Wzk, an ABC-transporter protein homologous to PglK from *Campylobacter jejuni*, and subsequently ligated to the lipid A-core by the Waal ligase (Hug *et al.*, 2010).

The presence of terminal fucosylated sugars on the outer surface of the bacterium, in particular the most common Le^X and Le^Y structures, is critical for colonization in mice models (Logan *et al.*, 2000, Moran *et al.*, 2000). However, the diversity of Lewis antigen expression in *H. pylori* hampers efforts to clearly define the role of these molecules in infection and disease progression. In humans, Le^X *H. pylori* O-specific oligosaccharide is recognized by galectin-3, a β -galactoside-binding lectin that serves as a gastric receptor (Fowler *et al.*, 2006). However, the main role attributed to the Lewis antigens is that of molecular mimicry, which could be manifested in several ways. For example, *H. pylori* can change its Lewis antigens in response to those present in the host, as demonstrated with Le^b-transgenic mice infected with Le^X-expressing *H. pylori*, which over time switched on Le^b expression (Pohl *et al.*, 2009). This change allowed better bacterial colonization than in the transgenic mice lacking Le^b expression, suggesting that Le^b *H. pylori* could survive better in a self-tolerant Le^b host (Pohl *et al.*, 2009). Alternatively, *H. pylori* expressing different Lewis antigens than those in the host can induce production of autoantibodies that recognize gastric parietal cells leading to disease (Negrini *et al.*, 1996, Faller *et al.*, 1997). Further, Lewis antigens can also dampen host immune responses to *H. pylori* through interactions with the C-type lectin DC-SIGN on

the surface of gastric dendritic cells, which lead to a block in maturation of T-helper 1 cells and reduced production of pro-inflammatory cytokines (Bergman *et al.*, 2004).

The first evidence that a single strain of *H. pylori* alters its LPS antigenic phenotype during the course of infection was demonstrated by investigating the expression of Lewis antigens in 127 isolates recovered from serial biopsies of 26 asymptomatic subjects (Rasko *et al.*, 2000). This alteration of LPS biosynthesis in *H. pylori* occurs during host colonization in response to several stimuli (Salaün *et al.*, 2005, Nilsson *et al.*, 2008) such as interaction with T helper cells (Bergman, 2004) and gastric pH (Skoglund *et al.*, 2009). More recently, several studies have focused on the genomic changes occurring in *H. pylori* isolates that have been recovered several years apart from patients with chronic infection (Falush *et al.*, 2001, Israel *et al.*, 2001, Kraft *et al.*, 2006, Alvi *et al.*, 2007, Morelli *et al.*, 2010, Kennemann *et al.*, 2011).

A whole-genome analysis of 10 *H. pylori* sequential isolates recovered from 4 patients over 16 years of chronic gastritis revealed 5 SNPs affecting LPS genes, including genes involved in the biosynthesis of lipid A (phosphoethanolamine transferase), core (*kdsA* and *waaF*) and O-specific oligosaccharide (*wecA*) synthesis, as well as in a putative lipopolysaccharide biosynthetic protein (Kennemann *et al.*, 2011). A cluster of nucleotide polymorphisms in the *fucT* (fucosyltransferase) gene, presumably facilitating its expression, was identified in whole-genome analyses of two *H. pylori* strains isolated from spouses (Linz *et al.*, 2013). Hyperexpression of *fucT* promotes posttranslational fucosylation of the O-specific oligosaccharide, generating Lewis antigens (Ge *et al.*, 1997, Martin *et al.*, 1997, Moran, 2008, Linz *et al.*, 2013). The alteration of *H. pylori* LPS during chronic gastric infection, either by altering LPS biosynthesis or by adding fucosyl residues to O-specific oligosaccharides, generates Lewis structures that mimic host antigens and contribute to immune system evasion.

MOLECULAR MECHANISMS OF LPS VARIATION

Antigenic variation of surface structures is a powerful mechanism for pathogen evasion of adaptive immune responses (Lerouge & Vanderleyden, 2002, van der Woude & Bäumlér, 2004, Lukáčová *et al.*, 2008). One of these adaptations involves phase variation, which is a

reversible, yet heritable form, of gene regulation that results in heterogeneous clonal populations and can be mediated by various molecular mechanisms (van der Woude & Bäumler, 2004). LPS phase variation can occur by addition of carbohydrates through the activity of glycosyltransferases or sialyltransferases, or addition of phosphorylcholine (ChoP) resulting in changes that affect antigenicity, serum sensitivity and adhesion (van der Woude & Bäumler, 2004). Phase variation has been described for human pathogens such as *S. enterica* serovar Typhimurium, *Campylobacter jejuni*, *Neisseria* spp. and *H. pylori* but because variable LPS modification is not easily identified, it is possible that phase variation is more widespread than currently known. Genetic and epigenetic mechanisms behind LPS variation are discussed below.

Adaptive mutagenesis and altered gene expression

Acquisition of adaptive mutations is a common theme in microbial persistence. In CF patients with chronic lung infection, *P. aeruginosa* strains accumulate a large proportion of mutator strains (Oliver *et al.*, 2000) that contribute to selection of mucoid variants (Oliver *et al.*, 2000, Mathee *et al.*, 2008, Ciofu *et al.*, 2010, Hogardt & Heesemann, 2010). The proinflammatory microenvironment in the airways including polymorphonuclear cells, hydrogen peroxide production, and antibiotics (Blázquez *et al.*, 2006) has been associated with mutagenesis and mucoid conversion *in vitro* (Mathee *et al.*, 1999, Sanders *et al.*, 2006, Moyano *et al.*, 2007). Cationic antimicrobial peptides can also exert a mutagenic inducing effect, as recently demonstrated for human cathelicidin LL-37 (Limoli *et al.*, 2014). Mutagenesis depended on LL-37 entering the bacterial cytosol and binding to DNA, which in turns promotes abnormal DNA synthesis by the error-prone polymerase DinB (Sanders *et al.*, 2006, Limoli *et al.*, 2014).

Environmental cues, such as ionic concentration, can lead to O antigen structural variations resulting from altered gene expression regulated by two-component signal transduction systems. One of the best examples of this type of regulation is the PhoP/PhoQ system in *Salmonella* (Prost & Miller, 2008, Needham & Trent, 2013). PhoQ is a membrane sensor histidine kinase and PhoP is its cognate response regulator. Activation of the PhoP/PhoQ system by acidic pH, specific antimicrobial peptides, and depletion of Mg^{2+} and Ca^{2+} stimulates transcription of *pagP* and *pagL* (among other genes) and subsequent

upregulation of the encoded proteins, which acylate and deacylate lipid A, respectively (Prost & Miller, 2008, Needham & Trent, 2013). Further, CF clinical isolates of *P. aeruginosa* obtained from patients treated with inhaled colistin (polymyxin E) can develop resistance by loss-of-function mutations in the *phoQ* gene (Miller *et al.*, 2011). Disruption of *phoQ* in the presence of an intact *phoP* stimulated Arap4N addition to lipid A by upregulated expression of the Arap4N synthesis operon. Therefore, this adaptive mutagenesis strategy results in high-level polymyxin resistance clinical strains of *P. aeruginosa*.

Slipped-strand mispairing

One of the molecular mechanisms of phase variation involves slipping of one of the DNA strands, which causes mispairing between daughter and parent strands during DNA replication (slipped-strand mispairing) (Lukáčová *et al.*, 2008). Short DNA repeats, microsatellites and tandem repeats are particularly prone to slipped-strand mispairing (van Belkum *et al.*, 1997, Torres-Cruz & van der Woude, 2003). In *H. pylori*, phase variation is related to an increase in the number of poly-C tract repeats in the β -(1,3)-galactosyl transferase (GalT), which leads to a switching on Le^b expression (Pohl *et al.*, 2009). Also, repetitive poly-A and poly-C sequences in the fucosyltransferase *fucT* mediate slipped-strand mispairing, which in turn results in production of Lewis antigens with different fucosylated oligosaccharides (Wang *et al.*, 2000, Nilsson *et al.*, 2008). Further, the α -(1,2)-fucosyltransferase gene *futC* contains an heptameric sequence (AAAAAAG) next to the ribosome binding site, which may cause a phase shift in the reading frame during translation (Wang *et al.*, 2000).

Lateral gene transfer, recombination, and genetic rearrangements

The heterogeneity of O antigens is mostly due to variation within the O antigen gene cluster, but it is unclear how such variation was generated (Reeves *et al.*, 2013). Genes involved in O antigen biosynthesis are generally arranged in large operons with low G+C content relative to the average G+C characteristic of each species, which suggests that these clusters were acquired by horizontal gene transfer from a species with low G+C content (Lerouge & Vanderleyden, 2002). The G+C content within the O antigen clusters also greatly differs from

gene to gene, indicating that the gene clusters might have been assembled from multiple horizontal transmission events and from several sources over a much longer time (Lerouge & Vanderleyden, 2002). The role of lateral gene transfer in the evolution of O antigen clusters and O antigen diversification has been well described in *Salmonella* (Perepelov *et al.*, 2011, Reeves *et al.*, 2013), *Escherichia* (D'Souza *et al.*, 2005, Hu *et al.*, 2010, Azmuda *et al.*, 2012), *Vibrio* (González-Fraga *et al.*, 2008, Wildschutte *et al.*, 2010), *Yersinia* (Cunneen & Reeves, 2007) and *Brucella* (Wattam *et al.*, 2014). Another mechanism of variation involves large chromosomal rearrangements. For example, more than half of the *P. aeruginosa* clone C isolates from CF lung infection exhibit large chromosomal inversions mediated an IS6100-induced coupled insertion-inversion mechanism. This creates also a selective advantage by insertion of IS6100 into *wbpM*, *pilB* and *mutS*, which leads to common CF phenotypes such as O-antigen and type IV pili deficiency and hyper mutability (Kresse *et al.*, 2003).

CONCLUDING REMARKS

The LPS is an abundant molecule of the outer membrane of most Gram-negative bacteria and plays a key role during host-pathogen interaction and the establishment of chronic infection. LPS-mediated virulence resides both in the endotoxic activity of lipid A and in the ability of the core and O antigen to provide the bacterium with resistance to host defence mechanisms. O antigen modification in general contributes to enhance the bacteria's ability to establish infection. For example, *P. aeruginosa* O antigen modification directed by the D3 prophage promotes adhesion to epithelial cells (Vaca-Pacheco *et al.*, 1999), while in *H. pylori*, expression of the Lewis antigen Le^X promotes bacterial adhesion to the gastric epithelia by interacting with host lectins. Further, O antigen modification can contribute to host immune evasion either by mimicry of host molecules (e.g., Lewis antigens in *H. pylori*) or by inhibiting activation of the host complement system (Raetz & Whitfield, 2002). It is also well established that during chronic infection there is an increase of mutator phenotypes (Oliver *et al.*, 2000), which leads to a higher mutation rate and will consequently contribute to the accumulation of modifications in LPS structure during colonization.

Several studies have shown alterations in the LPS molecule during chronic infection, which are thought to contribute to adhesion, host colonization, immune defenses evasion and

adaptation to the infection niche. Different mechanisms, both at the genetic and epigenetic levels have been implied in LPS variation, creating LPS diversity and thus contributing to the success of the infection.

Future progress in LPS research will require interdisciplinary experimental approaches, combining the application of genome-wide approaches (such as genomics, transcriptomics, proteomics and metabolomics), structural biology, animal knockout models, enzymology, carbohydrate chemistry and membrane biochemistry. LPS phase variation has been described for some human pathogens (*S. enterica* serovar Typhimurium, *C. jejuni*, *Neisseria* spp. and *H. pylori*) and future research should address the investigation of these mechanisms in other species as well. An in-depth understanding of LPS variation and its effects on pathogenicity and virulence is of paramount importance in the understanding of infection establishment and progression.

ACKNOWLEDGEMENTS

Research in the authors' laboratories has been supported the grants from the Canadian Institutes of Health Research, Cystic Fibrosis Canada, and the UK Cystic Fibrosis Trust (to M.A.V.) and from the Portuguese Foundation for Science and Technology (FCT) ((UID/BIO/04565/2013) and Programa Operacional Regional de Lisboa 2020 (Project N. 007317) to iBB - Institute for Bioengineering and Biosciences (to I.S-C.). R.F.M. was supported by the PhD fellowship SFRH/BD/84233/2012. I.S-C and M.A.V. were members of the EU COST Action BM1003: Microbial cell surface determinants of virulence as targets for new therapeutics in cystic fibrosis (<http://www.cost-bm1003.info/>) and acknowledge this support in the collation of this panel. The authors declare no conflict of interest.

REFERENCES

- Aaron SD, Vandemheen KL, Ramotar K, *et al.* (2010) Infection with transmissible strains of *Pseudomonas aeruginosa* and clinical outcomes in adults with cystic fibrosis. *JAMA* **304**: 2145-2153.
- Abeyrathne PD, Daniels C, Poon KK, Matewish MJ & Lam JS (2005) Functional characterization of WaaL, a ligase associated with linking O-antigen polysaccharide to the core of *Pseudomonas aeruginosa* lipopolysaccharide. *J Bacteriol* **187**: 3002-3012.
- Alvi A, Devi SM, Ahmed I, *et al.* (2007) Microevolution of *Helicobacter pylori* type IV secretion systems in an ulcer disease patient over a ten-year period. *J Clin Microbiol* **45**: 4039-4043.
- Aspinall GO, Monteiro MA, Pang H, Walsh EJ & Moran AP (1996) Lipopolysaccharide of the *Helicobacter pylori* type strain NCTC 11637 (ATCC 43504): structure of the O antigen chain and core oligosaccharide regions. *Biochemistry* **35**: 2489-2497.
- Azmuda N, Rahman MZ, Sultana M, Jenssen EL, Khan SI & Birkeland N-K (2012) Evidence of interspecies O antigen gene cluster transfer between *Shigella boydii* 15 and *Escherichia fergusonii*. *APMIS* **120**: 959-966.
- Babu M, Diaz-Mejia JJ, Vlasblom J, *et al.* (2011) Genetic interaction maps in *Escherichia coli* reveal functional crosstalk among cell envelope biogenesis pathways. *PLoS Genet* **7**: e1002377.
- Baker PJ, Hraba T, Taylor CE, *et al.* (1994) Molecular structures that influence the immunomodulatory properties of the lipid A and inner core region oligosaccharides of bacterial lipopolysaccharides. *Infect Immun* **62**: 2257-2269.
- Berg DE, Hoffman PS, Appelmek BJ & Kusters JG (1997) The *Helicobacter pylori* genome sequence: genetic factors for long life in the gastric mucosa. *Trends Microbiol* **5**: 468-474.
- Bergman MP, Engering A, Smits HH, *et al.* (2004) *Helicobacter pylori* modulates the T helper cell 1/T helper cell 2 balance through phase-variable interaction between lipopolysaccharide and DC-SIGN. *J Exp Med* **200**: 979-990.
- Berry MC, McGhee GC, Zhao Y & Sundin GW (2009) Effect of a *waaL* mutation on lipopolysaccharide composition, oxidative stress survival, and virulence in *Erwinia amylovora*. *FEMS Microbiol Lett* **291**: 80-87.
- Blázquez J, Gómez-Gómez JM, Oliver A, Juan C, Kapur V & Martín S (2006) PBP3 inhibition elicits adaptive responses in *Pseudomonas aeruginosa*. *Mol Microbiol* **62**: 84-99.
- Bowden SD, Hale N, Chung JC, Hodgkinson JT, Spring DR & Welch M (2013) Surface swarming motility by *Pectobacterium atrosepticum* is a latent phenotype that requires O antigen and is regulated by quorum sensing. *Microbiology* **159**: 2375-2385.
- Bragonzi A, Wiehlmann L, Klockgether J, Cramer N, Worlitzsch D, Döring G & Tümmler B (2006) Sequence diversity of the *mucABD* locus in *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *Microbiology* **152**: 3261-3269.
- Brown LM (2000) *Helicobacter pylori*: epidemiology and routes of transmission. *Epidemiol Rev* **22**: 283-297.
- Bystrova OV, Knirel YA, Lindner B, Kocharova NA, Kondavova AN, Zähringer U & Pier GB (2006) Structures of the core oligosaccharide and O-units in the R- and SR-type lipopolysaccharides of reference strains of *Pseudomonas aeruginosa* O-serogroups. *FEMS Immunol Med Microbiol* **46**: 85-99.

635 Chmiela M, Mischczyk E & Rudnicka K (2014) Structural modifications of *Helicobacter pylori*
636 lipopolysaccharide: an idea for how to live in peace. *World J Gastroenterol* **20**: 9882-
637 9897.

638 Chung HS & Raetz CR (2011) Dioxygenases in *Burkholderia ambifaria* and *Yersinia pestis*
639 that hydroxylate the outer Kdo unit of lipopolysaccharide. *Proc Natl Acad Sci U S A* **108**:
640 510-515.

641 Cigana C, Curcurù L, Leone MR, *et al.* (2009) *Pseudomonas aeruginosa* exploits lipid A and
642 mucopeptides modification as a strategy to lower innate immunity during cystic fibrosis
643 lung infection. *PLoS one* **4**: e8439.

644 Ciofu O, Mandsberg LF, Bjarnsholt T, Wassermann T & Høiby N (2010) Genetic adaptation of
645 *Pseudomonas aeruginosa* during chronic lung infection of patients with cystic fibrosis:
646 strong and weak mutators with heterogeneous genetic backgrounds emerge in *mucA*
647 and/or *lasR* mutants. *Microbiology* **156**: 1108-1119.

648 Ciofu O, Tolker-Nielsen T, Jensen PO, Wang H & Høiby N (2015) Antimicrobial resistance,
649 respiratory tract infections and role of biofilms in lung infections in cystic fibrosis patients.
650 *Adv Drug Deliv Rev* **85**: 7-23.

651 Coutinho CP, de Carvalho CC, Madeira A, Pinto-de-Oliveira A & Sa-Correia I (2011)
652 *Burkholderia cenocepacia* phenotypic clonal variation during a 3.5-year colonization in the
653 lungs of a cystic fibrosis patient. *Infect Immun* **79**: 2950-2960.

654 Coutinho CP, Dos Santos SC, Madeira A, Mira NP, Moreira AS & Sá-Correia I (2011) Long-
655 Term Colonization of the Cystic Fibrosis Lung by *Burkholderia cepacia* Complex Bacteria:
656 Epidemiology, Clonal Variation, and Genome-Wide Expression Alterations. *Front Cell*
657 *Infect Microbiol* **1**: 12.

658 Cramer N, Klockgether J, Wrasman K, Schmidt MG, Davenport CF & Tümmler B (2011)
659 Microevolution of the major common *Pseudomonas aeruginosa* clones C and PA14 in
660 cystic fibrosis lungs. *Environ Microbiol* **13**: 1690-1704.

661 Cullen L & McClean S (2015) Bacterial Adaptation during Chronic Respiratory Infections.
662 *Pathogens* **4**: 66-89.

663 Cullen TW, Giles DK, Wolf LN, Ecobichon C, Boneca IG & Trent MS (2011) *Helicobacter*
664 *pylori* versus the host: remodeling of the bacterial outer membrane is required for survival
665 in the gastric mucosa. *PLoS Pathogens* **7**: e1002454.

666 Cunneen MM & Reeves PR (2007) The *Yersinia kristensenii* O11 O-antigen gene cluster was
667 acquired by lateral gene transfer and incorporated at a novel chromosomal locus. *Mol*
668 *Biol Evol* **24**: 1355-1365.

669 D'Argenio DA, Wu M, Hoffman LR, *et al.* (2007) Growth phenotypes of *Pseudomonas*
670 *aeruginosa lasR* mutants adapted to the airways of cystic fibrosis patients. *Mol Microbiol*
671 **64**: 512-533.

672 D'Souza JM, Samuel GN & Reeves PR (2005) Evolutionary origins and sequence of the
673 *Escherichia coli* O4 O-antigen gene cluster. *FEMS Microbiol Lett* **244**: 27-32.

674 Darveau RP, Cunningham MD, Bailey T, *et al.* (1995) Ability of bacteria associated with
675 chronic inflammatory disease to stimulate E-selectin expression and promote neutrophil
676 adhesion. *Infect Immun* **63**: 1311-1317.

677 de Bernard M & Josenhans C (2014) Pathogenesis of *Helicobacter pylori* infection.
678 *Helicobacter* **19**: 11-18.

679 Delucia AM, Six DA, Caughlan RE, Gee P, Hunt I, Lam JS & Dean CR (2011)
680 Lipopolysaccharide (LPS) inner-core phosphates are required for complete LPS synthesis
681 and transport to the outer membrane in *Pseudomonas aeruginosa* PAO1. *MBio*
682 **Aug2;2(4)**.

683 Dettman JR, Rodrigue N, Aaron SD & Kassen R (2013) Evolutionary genomics of epidemic
684 and nonepidemic strains of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* **110**:
685 21065-21070.

686 Di Lorenzo F, Silipo A, Bianconi I, *et al.* (2015) Persistent cystic fibrosis isolate *Pseudomonas*
687 *aeruginosa* strain RP73 exhibits an under-acylated LPS structure responsible of its low
688 inflammatory activity. *Mol Immunol* **63**: 166-175.

689 Di Lorenzo F, Kubik L, Oblak A, *et al.* (2015) Activation of Human Toll-like Receptor 4
690 (TLR4).Myeloid Differentiation Factor 2 (MD-2) by Hypoacylated Lipopolysaccharide from
691 a Clinical Isolate of *Burkholderia cenocepacia*. *J Biol Chem* **290**: 21305-21319.

692 Drevinek P & Mahenthiralingam E (2010) *Burkholderia cenocepacia* in cystic fibrosis:
693 epidemiology and molecular mechanisms of virulence. *Clin. Microbiol. Infect.* **16**: 821-
694 830.

695 Duerr CU, Zenk SF, Chassin C, Pott J, Gutle D, Hensel M & Hornef MW (2009) O-antigen
696 delays lipopolysaccharide recognition and impairs antibacterial host defense in murine
697 intestinal epithelial cells. *PLoS Pathog.* **5**: e1000567.

698 Ernst RK, Hajjar AM, Tsai JH, Moskowitz SM, Wilson CB & Miller SI (2003) *Pseudomonas*
699 *aeruginosa* lipid A diversity and its recognition by Toll-like receptor 4. *J Endotoxin Res* **9**:
700 395-400.

701 Ernst RK, Adams KN, Moskowitz SM, *et al.* (2006) The *Pseudomonas aeruginosa* lipid A
702 deacylase: selection for expression and loss within the cystic fibrosis airway. *J Bacteriol*
703 **188**: 191-201.

704 Ernst RK, Moskowitz SM, Emerson JC, *et al.* (2007) Unique lipid a modifications in
705 *Pseudomonas aeruginosa* isolated from the airways of patients with cystic fibrosis. *J*
706 *Infect Dis* **196**: 1088-1092.

707 Faller G, Steininger H, Kränzlein J, *et al.* (1997) Antigastric autoantibodies in *Helicobacter*
708 *pylori* infection: implications of histological and clinical parameters of gastritis. *Gut* **41**:
709 619-623.

710 Falush D, Kraft C, Taylor NS, Correa P, Fox JG, Achtman M & Suerbaum S (2001)
711 Recombination and mutation during long-term gastric colonization by *Helicobacter pylori*:
712 estimates of clock rates, recombination size, and minimal age. *Proc Natl Acad Sci U S A*
713 **98**: 15056-15061.

714 Fowler M, Thomas RJ, Atherton J, Roberts IS & High NJ (2006) Galectin-3 binds to
715 *Helicobacter pylori* O-antigen: it is upregulated and rapidly secreted by gastric epithelial
716 cells in response to *H. pylori* adhesion. *Cell Microbiol* **8**: 44-54.

717 Ge Z, Chan NW, Palcic MM & Taylor DE (1997) Cloning and heterologous expression of an
718 α 1,3-fucosyltransferase gene from the gastric pathogen *Helicobacter pylori*. *J Biol Chem*
719 **272**: 21357-21363.

720 Geurtsen J, Steeghs L, Hove JT, van der Ley P & Tommassen J (2005) Dissemination of lipid
721 A deacylases (*pagL*) among Gram-negative bacteria: identification of active-site histidine
722 and serine residues. *J Biol Chem* **280**: 8248-8259.

- 723 González-Fraga S, Pichel M, Binsztein N, Johnson JA, Morris JG & Stine OC (2008) Lateral
724 gene transfer of O1 serogroup encoding genes of *Vibrio cholerae*. *FEMS Microbiol Lett*
725 **286**: 32-38.
- 726 Govan JRW & Deretic V (1996) Microbial pathogenesis in cystic fibrosis: mucoid
727 *Pseudomonas aeruginosa* and *Burkholderia cepacia*. *Microbiol Rev* **60**: 539-574.
- 728 Greenfield LK & Whitfield C (2012) Synthesis of lipopolysaccharide O-antigens by ABC
729 transporter-dependent pathways. *Carbohydr Res* **356**: 12-24.
- 730 Hancock RE, Mutharia LM, Chan L, Darveau RP, Speert DP & Pier GB (1983) *Pseudomonas*
731 *aeruginosa* isolates from patients with cystic fibrosis: a class of serum-sensitive,
732 nontypable strains deficient in lipopolysaccharide O side chains. *Infect Immun* **42**: 170-
733 177.
- 734 Hao Y, King JD, Huszczyński S, Kocincova D & Lam JS (2013) Five new genes are important
735 for common polysaccharide antigen biosynthesis in *Pseudomonas aeruginosa*. *MBio* **4**:
736 e00631-00612.
- 737 Hatakeyama M (2014) *Helicobacter pylori* CagA and gastric cancer: a paradigm for hit-and-
738 run carcinogenesis. *Cell Host and Microbe* **15**: 306-316.
- 739 Hauser AR, Jain M, Bar-Meir M & McColley SA (2011) Clinical significance of microbial
740 infection and adaptation in cystic fibrosis. *Clin Microbiol Rev* **24**: 29-70.
- 741 Hoffman LR, Kulasekara HD, Emerson JC, Houston LS, Burns JL, Ramsey BW & Miller SI
742 (2009) *Pseudomonas aeruginosa lasR* mutants are associated with cystic fibrosis lung
743 disease progression. *J Cyst Fibros* **8**: 66-70.
- 744 Hogardt M & Heesemann J (2010) Adaptation of *Pseudomonas aeruginosa* during
745 persistence in the cystic fibrosis lung. *Int J Med Microbiol* **300**: 557-562.
- 746 Hu B, Perepelov AV, Liu B, *et al.* (2010) Structural and genetic evidence for the close
747 relationship between *Escherichia coli* O71 and *Salmonella enterica* O28 O-antigens.
748 *FEMS Immunol Med Microbiol* **59**: 161-169.
- 749 Hug I, Couturier MR, Rooker MM, Taylor DE, Stein M & Feldman MF (2010) *Helicobacter*
750 *pylori* lipopolysaccharide is synthesized via a novel pathway with an evolutionary
751 connection to protein N-glycosylation. *PLoS Path.* **6**: e1000819.
- 752 Israel DA, Salama N, Krishna U, Rieger UM, Atherton JC, Falkow S & Peek RM (2001)
753 *Helicobacter pylori* genetic diversity within the gastric niche of a single human host. *Proc.*
754 *Natl. Acad. Sci. USA* **98**: 14625-14630.
- 755 Jain M, Ramirez D, Seshadri R, *et al.* (2004) Type III secretion phenotypes of *Pseudomonas*
756 *aeruginosa* strains change during infection of individuals with cystic fibrosis. *J Clin*
757 *Microbiol* **42**: 5229-5237.
- 758 Kadurugamuwa JL, Lam JS & Beveridge TJ (1993) Interaction of gentamicin with the A band
759 and B band lipopolysaccharides of *Pseudomonas aeruginosa* and its possible lethal
760 effect. *Antimicrob Agents Chemother* **37**: 715-721.
- 761 Kang J & Blaser MJ (2006) Bacterial populations as perfect gases: genomic integrity and
762 diversification tensions in *Helicobacter pylori*. *Nature Rev Microbiol* **4**: 826-836.
- 763 Keenleyside WJ & Whitfield C (1996) A novel pathway for O-polysaccharide biosynthesis in
764 *Salmonella enterica* serovar Borreze. *J. Biol. Chem.* **271**: 28581-28592.
- 765 Kennemann L, Didelot X, Aebischer T, *et al.* (2011) *Helicobacter pylori* genome evolution
766 during human infection. *Proc Natl Acad Sci U S A* **108**: 5033-5038.

767 King JD, Kocincova D, Westman EL & Lam JS (2009) Review: Lipopolysaccharide
768 biosynthesis in *Pseudomonas aeruginosa*. *Innate Immun* **15**: 261-312.

769 Knirel YA, Bystrova OV, Kocharova NA, Zahringer U & Pier GB (2006) Conserved and
770 variable structural features in the lipopolysaccharide of *Pseudomonas aeruginosa*. *J*
771 *Endotoxin Res* **12**: 324-336.

772 Kraft C, Stack A, Josenhans C, *et al.* (2006) Genomic changes during chronic *Helicobacter*
773 *pylori* infection. *J Bacteriol* **188**: 249-254.

774 Kresse AU, Dinesh SD, Larbig K & Römling U (2003) Impact of large chromosomal inversions
775 on the adaptation and evolution of *Pseudomonas aeruginosa* chronically colonizing cystic
776 fibrosis lungs. *Mol. Microbiol.* **47**: 145-158.

777 Lam JS, Taylor VL, Islam ST, Hao Y & Kocincova D (2011) Genetic and Functional Diversity
778 of *Pseudomonas aeruginosa* Lipopolysaccharide. *Front Microbiol* **2**: 118.

779 Lee B, Haagensen JAJ, Ciofu O, Andersen JB, Høiby N & Molin S (2005) Heterogeneity of
780 biofilms formed by nonmucoid *Pseudomonas aeruginosa* isolates from patients with
781 cystic fibrosis. *J Clin Microbiol* **43**: 5247-5255.

782 Lerouge I & Vanderleyden J (2002) O-antigen structural variation: mechanisms and possible
783 roles in animal/plant-microbe interactions. *FEMS Microbiol Rev* **26**: 17-47.

784 Lieberman TD, Michel J-B, Aingaran M, *et al.* (2011) Parallel bacterial evolution within
785 multiple patients identifies candidate pathogenicity genes. *Nat Genet* **43**: 1275-1280.

786 Limoli DH, Rockel AB, Host KM, Jha A, Kopp BT, Hollis T & Wozniak DJ (2014) Cationic
787 antimicrobial peptides promote microbial mutagenesis and pathoadaptation in chronic
788 infections. *PLoS Pathogens* **10**: e1004083.

789 Linz B, Windsor HM, Gajewski JP, Hake CM, Drautz DI, Schuster SC & Marshall BJ (2013)
790 *Helicobacter pylori* genomic microevolution during naturally occurring transmission
791 between adults. *Plos One* **8**: e82187.

792 Lipuma JJ (2010) The changing microbial epidemiology in cystic fibrosis. *Clin Microbiol Rev*
793 **23**: 299-323.

794 Ljungh Å, Moran AP & Wadström T (1996) Interactions of bacterial adhesins with extracellular
795 matrix and plasma proteins: pathogenic implications and therapeutic possibilities. *FEMS*
796 *Immunol Med Microbiol* **16**: 117-126.

797 Logan SM, Conlan JW, Monteiro Ma, Wakarchuk WW & Altman E (2000) Functional
798 genomics of *Helicobacter pylori*: identification of a β -1,4 galactosyltransferase and
799 generation of mutants with altered lipopolysaccharide. *Mol Microbiol* **35**: 1156-1167.

800 Lorè NI, Cigana C, De Fino I, *et al.* (2012) Cystic fibrosis-niche adaptation of *Pseudomonas*
801 *aeruginosa* reduces virulence in multiple infection hosts. *PloS one* **7**: e35648.

802 Loutet SA & Valvano MA (2010) A decade of *Burkholderia cenocepacia* virulence determinant
803 research. *Infect Immun* **78**: 4088-4100.

804 Lukáčová M, Barák I & Kazár J (2008) Role of structural variations of polysaccharide antigens
805 in the pathogenicity of Gram-negative bacteria. *Clin Microbiol Infect* **14**: 200-206.

806 Lyczak JB, Cannon CL & Pier GB (2002) Lung infections associated with cystic fibrosis. *Clin.*
807 *Microbiol. Rev.* **15**: 194-222.

- 808 Madeira A, Santos PM, Coutinho CP, Pinto-de-Oliveira A & Sa-Correia I (2011) Quantitative
809 proteomics (2-D DIGE) reveals molecular strategies employed by *Burkholderia*
810 *cenocepacia* to adapt to the airways of cystic fibrosis patients under antimicrobial therapy.
811 *Proteomics* **11**: 1313-1328.
- 812 Madeira A, dos Santos SC, Santos PM, *et al.* (2013) Proteomic profiling of *Burkholderia*
813 *cenocepacia* clonal isolates with different virulence potential retrieved from a cystic
814 fibrosis patient during chronic lung infection. *PLoS One* **8**: e83065.
- 815 Mahenthiralingam E, Campbell ME & Speert DP (1994) Nonmotility and phagocytic resistance
816 of *Pseudomonas aeruginosa* isolates from chronically colonized patients with cystic
817 fibrosis. *Infection & Immunity* **62**: 596-605.
- 818 Mahenthiralingam E, Urban TA & Goldberg JB (2005) The multifarious, multireplicon
819 *Burkholderia cepacia* complex. *Nat. Rev. Microbiol.* **3**: 144-156.
- 820 Malnick SD, Melzer E, Attali M, Duek G & Yahav J (2014) *Helicobacter pylori*: friend or foe?
821 *World J Gastroenterol* **20**: 8979-8985.
- 822 Mamat U, Skurnik M & Bengoechea JA (2011) LPS core oligosaccharide biosynthesis and
823 assembly. *Bacterial lipopolysaccharides: Structure, chemical synthesis, biogenesis and*
824 *interaction with host cells*, (Knirel YA & Valvano MA, ed.^eds.), p.^pp. 237-273. Springer-
825 Verlag, Wien.
- 826 Martin SL, Edbrooke MR, Hodgman TC, van den Eijnden DH & Bird MI (1997) Lewis X
827 biosynthesis in *Helicobacter pylori*. Molecular cloning of an $\alpha(1,3)$ -fucosyltransferase
828 gene. *J Biol Chem* **272**: 21349-21356.
- 829 Mathee K, Ciofu O, Sternberg C, *et al.* (1999) Mucoïd conversion of *Pseudomonas*
830 *aeruginosa* by hydrogen peroxide: a mechanism for virulence activation in the cystic
831 fibrosis lung. *Microbiology* **145**: 1349-1357.
- 832 Mathee K, Narasimhan G, Valdes C, *et al.* (2008) Dynamics of *Pseudomonas aeruginosa*
833 genome evolution. *Proc Natl Acad Sci U S A* **105**: 3100-3105.
- 834 Mattsby-Baltzer I, Mielniczuk Z, Larsson L, Lindgren K & Goodwin S (1992) Lipid A in
835 *Helicobacter pylori*. *Infect Immun* **60**: 4383-4387.
- 836 May JM, Sherman DJ, Simpson BW, Ruiz N & Kahne D (2015) Lipopolysaccharide transport
837 to the cell surface: periplasmic transport and assembly into the outer membrane. *Philos*
838 *Trans R Soc Lond B Biol Sci* **370**.
- 839 Mégraud F, Bessède E & Varon C (2015) *Helicobacter pylori* infection and gastric carcinoma.
840 *Clin Microbiol Infect* **9**: 222-223.
- 841 Miller AK, Brannon MK, Stevens L, *et al.* (2011) PhoQ mutations promote lipid A modification
842 and polymyxin resistance of *Pseudomonas aeruginosa* found in colistin-treated cystic
843 fibrosis patients. *Antimicrob Agents Chemother* **55**: 5761-5769.
- 844 Mira NP, Madeira A, Moreira AS, Coutinho CP & Sá-Correia I (2011) Genomic expression
845 analysis reveals strategies of *Burkholderia cenocepacia* to adapt to cystic fibrosis
846 patients' airways and antimicrobial therapy. *PloS one* **6**: e28831.
- 847 Monteiro MA, Chan KH, Rasko DA, *et al.* (1998) Simultaneous expression of type 1 and type
848 2 Lewis blood group antigens by *Helicobacter pylori* lipopolysaccharides. Molecular
849 mimicry between *H. pylori* lipopolysaccharides and human gastric epithelial cell surface
850 glycoforms. *J Biol Chem* **273**: 11533-11543.
- 851 Moran AP (2008) Relevance of fucosylation and Lewis antigen expression in the bacterial
852 gastroduodenal pathogen *Helicobacter pylori*. *Carbohydr Res* **343**: 1952-1965.

853 Moran AP & Aspinall GO (1998) Unique structural and biological features of *Helicobacter*
854 *pylori* lipopolysaccharides. *Prog Clin Biol Res* **397**: 37-49.

855 Moran AP, Prendergast MM & Appelmek BJ (1996) Molecular mimicry of host structures by
856 bacterial lipopolysaccharides and its contribution to disease. *FEMS Immunol Med*
857 *Microbiol* **16**: 105-115.

858 Moran AP, Lindner B & Walsh EJ (1997) Structural characterization of the lipid A component
859 of *Helicobacter pylori* rough- and smooth-form lipopolysaccharides. *J Bacteriol* **179**: 6453-
860 6463.

861 Moran AP, Sturegård E, Sjunnesson H, Wadström T & Hynes SO (2000) The relationship
862 between O-chain expression and colonisation ability of *Helicobacter pylori* in a mouse
863 model. *FEMS Immunol Med Microbiol* **29**: 263-270.

864 Moreira AS, Coutinho CP, Azevedo P, Lito L, Melo-Cristino J & Sá-Correia I (2014)
865 *Burkholderia dolosa* phenotypic variation during the decline in lung function of a cystic
866 fibrosis patient during 5.5 years of chronic colonization. *J Med Microbiol* **63**: 594-601.

867 Morelli G, Didelot X, Kusecek B, *et al.* (2010) Microevolution of *Helicobacter pylori* during
868 prolonged infection of single hosts and within families. *PLoS genetics* **6**: e1001036.

869 Moskowitz SM & Ernst RK (2010) The role of *Pseudomonas lipopolysaccharide* in cystic
870 fibrosis airway infection. *Subcell Biochem* **53**: 241-253.

871 Moskowitz SM, Ernst RK & Miller SI (2004) PmrAB, a two-component regulatory system of
872 *Pseudomonas aeruginosa* that modulates resistance to cationic antimicrobial peptides
873 and addition of aminoarabinose to lipid A. *J. Bacteriol.* **186**: 575-579.

874 Mowat E, Paterson S, Fothergill JL, *et al.* (2011) *Pseudomonas aeruginosa* population
875 diversity and turnover in cystic fibrosis chronic infections. *Am J Respir Crit Care Med* **183**:
876 1674-1679.

877 Moyano AJ, Luján AM, Argaraña CE & Smania AM (2007) MutS deficiency and activity of the
878 error-prone DNA polymerase IV are crucial for determining *mucA* as the main target for
879 mucoid conversion in *Pseudomonas aeruginosa*. *Mol Microbiol* **64**: 547-559.

880 Muotiala A, Helander IM, Pyhala L, Kosunen TU & Moran AP (1992) Low biological activity of
881 *Helicobacter pylori* lipopolysaccharide. *Infect Immun* **60**: 1714-1716.

882 Murray GL, Attridge SR & Morona R (2006) Altering the length of the lipopolysaccharide O
883 antigen has an impact on the interaction of *Salmonella enterica* serovar Typhimurium with
884 macrophages and complement. *J. Bacteriol.* **188**: 2735-2739.

885 Needham BD & Trent MS (2013) Fortifying the barrier: the impact of lipid A remodelling on
886 bacterial pathogenesis. *Nat Rev Microbiol* **11**: 467-481.

887 Negrini R, Savio A, Poesi C, *et al.* (1996) Antigenic mimicry between *Helicobacter pylori* and
888 gastric mucosa in the pathogenesis of body atrophic gastritis. *Gastroenterology* **111**: 655-
889 665.

890 Nikaido H (2003) Molecular basis of bacterial outer membrane permeability revisited.
891 *Microbiol. Mol. Biol. Rev.* **67**: 593-656.

892 Nilsson C, Skoglund A, Moran AP, Annuk H, Engstrand L & Normark S (2008)
893 Lipopolysaccharide diversity evolving in *Helicobacter pylori* communities through genetic
894 modifications in fucosyltransferases. *Plos One* **3**: e3811.

895 Nowicki EM, O'Brien JP, Brodbelt JS & Trent MS (2015) Extracellular zinc induces
896 phosphoethanolamine addition to *Pseudomonas aeruginosa* lipid A via the ColRS two-
897 component system. *Mol Microbiol* **97**: 166-178.

898 Oberhardt MA, Goldberg JB, Hogardt M & Papin JA (2010) Metabolic network analysis of
899 *Pseudomonas aeruginosa* during chronic cystic fibrosis lung infection. *J Bacteriol* **192**:
900 5534-5548.

901 Oliver A, Canton R, Campo P, Baquero F & Blazquez J (2000) High frequency of
902 hypermutable *Pseudomonas aeruginosa* in cystic fibrosis lung infection. *Science* **288**:
903 1251-1254.

904 Otero LL, Ruiz VE & Perez Perez GI (2014) *Helicobacter pylori*: the balance between a role
905 as colonizer and pathogen. *Best Pract Res Clin Gastroenterol* **28**: 1017-1029.

906 Paixão TA, Roux CM, den Hartigh AB, Sankaran-Walters S, Dandekar S, Santos RL & Tsolis
907 RM (2009) Establishment of systemic *Brucella melitensis* infection through the digestive
908 tract requires urease, the type IV secretion system, and lipopolysaccharide O antigen.
909 *Infect. Immun.* **77**: 4197-4208.

910 Park BS & Lee JO (2013) Recognition of lipopolysaccharide pattern by TLR4 complexes. *Exp*
911 *Mol Med* **45**: e66.

912 Park BS, Song DH, Kim HM, Choi BS, Lee H & Lee JO (2009) The structural basis of
913 lipopolysaccharide recognition by the TLR4-MD-2 complex. *Nature* **458**: 1191-1195.

914 Parkins MD & Floto RA (2015) Emerging bacterial pathogens and changing concepts of
915 bacterial pathogenesis in cystic fibrosis. *J Cyst Fibros* **14**: 293-304.

916 Perepelov AV, Liu B, Guo D, *et al.* (2011) Structure elucidation of the O-Antigen of
917 *Salmonella enterica* O51 and its structural and genetic relation to the O-Antigen of
918 *Escherichia coli* O23. *Biochemistry* **76**: 774-779.

919 Pérez-Pérez GI, Shepherd VL, Morrow JD & Blaser MJ (1995) Activation of human THP-1
920 cells and rat bone marrow-derived macrophages by *Helicobacter pylori*
921 lipopolysaccharide. *Infect Immun* **63**: 1183-1187.

922 Pohl MA, Romero-Gallo J, Guruge JL, Tse DB, Gordon JI & Blaser MJ (2009) Host-
923 dependent Lewis (Le) antigen expression in *Helicobacter pylori* cells recovered from Leb-
924 transgenic mice. *J Exp Med* **206**: 3061-3072.

925 Poole K (2005) Efflux-mediated antimicrobial resistance. *J Antimicrob Chemother* **56**: 20-51.

926 Posselt G, Backert S & Wessler S (2013) The functional interplay of *Helicobacter pylori*
927 factors with gastric epithelial cells induces a multi-step process in pathogenesis. *Cell*
928 *Commun Signal* **11**: 77.

929 Prost LR & Miller SI (2008) The *Salmonellae* PhoQ sensor: mechanisms of detection of
930 phagosome signals. *Cell Microbiol* **10**: 576-582.

931 Raetz CR, Reynolds CM, Trent MS & Bishop RE (2007) Lipid A modification systems in gram-
932 negative bacteria. *Annu. Rev. Biochem.* **76**: 295-329.

933 Raetz CRH & Whitfield C (2002) Lipopolysaccharide endotoxins. *Annu. Rev. Biochem.* **71**:
934 635-700.

935 Rasko DA, Wilson TJ, Zopf D & Taylor DE (2000) Lewis antigen expression and stability in
936 *Helicobacter pylori* isolated from serial gastric biopsies. *J Infect Dis* **181**: 1089-1095.

937 Reeves PR, Cunneen MM, Liu B & Wang L (2013) Genetics and evolution of the *Salmonella*
938 galactose-initiated set of O antigens. *PloS One* **8**: e69306.

939 Reeves PR, Hobbs M, Valvano MA, *et al.* (1996) Bacterial polysaccharide synthesis and gene
940 nomenclature. *Trends Microbiol.* **4**: 495-503.

941 Ruan X, Pérez JM, Marolda CL & Valvano MA (2012) The WaaL O-antigen
942 lipopolysaccharide ligase has features in common with metal ion-independent inverting
943 glycosyltransferases. *Glycobiology* **22**: 288-299.

944 Rubin EJ & Trent MS (2013) Colonize, evade, flourish: how glyco-conjugates promote
945 virulence of *Helicobacter pylori*. *Gut Microbes* **4**: 439-453.

946 Sadovskaya I, Brisson JR, Thibault P, Richards JC, Lam JS & Altman E (2000) Structural
947 characterization of the outer core and the O-chain linkage region of lipopolysaccharide
948 from *Pseudomonas aeruginosa* serotype O5. *Eur. J. Biochem.* **267**: 1640-1650.

949 Salaün L, Ayraud S & Saunders NJ (2005) Phase variation mediated niche adaptation during
950 prolonged experimental murine infection with *Helicobacter pylori*. *Microbiology* **151**: 917-
951 923.

952 Saldías MS, Ortega X & Valvano MA (2009) *Burkholderia cenocepacia* O antigen
953 lipopolysaccharide prevents phagocytosis by macrophages and adhesion to epithelial
954 cells. *J. Med. Microbiol.* **58**: 1542-1548.

955 Sanders LH, Rockel A, Lu H, Wozniak DJ & Sutton MD (2006) Role of *Pseudomonas*
956 *aeruginosa* *dinB*-encoded DNA polymerase IV in mutagenesis. *J Bacteriol* **188**: 8573-
957 8585.

958 Schroeder TH, Lee MM, Yacono PW, Cannon CL, Gerçeker AA, Golan DE & Pier GB (2002)
959 CFTR is a pattern recognition molecule that extracts *Pseudomonas aeruginosa* LPS from
960 the outer membrane into epithelial cells and activates NF-kappa B translocation. *Proc.*
961 *Natl. Acad. Sci. U S A* **99**: 6907-6912.

962 Schwab U, Abdullah LH, Perlmutter OS, *et al.* (2014) Localization of *Burkholderia cepacia*
963 complex bacteria in cystic fibrosis lungs and interactions with *Pseudomonas aeruginosa*
964 in hypoxic mucus. *Infect Immun* **82**: 4729-4745.

965 Semeraro N, Montemurro P, Piccoli C, *et al.* (1996) Effect of *Helicobacter pylori*
966 lipopolysaccharide (LPS) and LPS derivatives on the production of tissue factor and
967 plasminogen activator inhibitor type 2 by human blood mononuclear cells. *J Infect Dis*
968 **174**: 1255-1260.

969 Shapiro AB, Gu RF & Gao N (2014) Dimerization of isolated *Pseudomonas aeruginosa*
970 lipopolysaccharide transporter component LptA. *Biochem Biophys Res Commun* **450**:
971 1327-1332.

972 Silipo A, Molinaro A, Cescutti P, Bedini E, Rizzo R, Parrilli M & Lanzetta R (2005) Complete
973 structural characterization of the lipid A fraction of a clinical strain of *B. cepacia*
974 genomovar I lipopolysaccharide. *Glycobiology* **15**: 561-570.

975 Silipo A, Molinaro A, Ierano T, *et al.* (2007) The complete structure and pro-inflammatory
976 activity of the lipooligosaccharide of the highly epidemic and virulent Gram-negative
977 bacterium *Burkholderia cenocepacia* ET-12 (strain J2315). *Chemistry* **13**: 3501-3511.

978 Simpson BW, May JM, Sherman DJ, Kahne D & Ruiz N (2015) Lipopolysaccharide transport
979 to the cell surface: biosynthesis and extraction from the inner membrane. *Philos Trans R*
980 *Soc Lond B Biol Sci* **370**.

981 Skoglund A, Bäckhed HK, Nilsson C, Bjorkholm B, Normark S & Engstrand L (2009) A
982 changing gastric environment leads to adaptation of lipopolysaccharide variants in
983 *Helicobacter pylori* populations during colonization. *PLoS One* **4**: e5885.

984 Smith EE, Buckley DG, Wu Z, *et al.* (2006) Genetic adaptation by *Pseudomonas aeruginosa*
985 to the airways of cystic fibrosis patients. *Proc Natl Acad Sci U S A* **103**: 8487-8492.

986 Sperandio P, Dehò G & Polissi A (2011) Lipopolysaccharide export to the outer membrane.
987 *Bacterial lipopolysaccharides: Structure, Chemical synthesis, biogenesis and interaction*
988 *with host cells*, (Knirel Y & Valvano MA, ed.[^]eds.), p.[^]pp. 311-337. Springer-Verlag, Wien.

989 Sperandio P, Villa R, Martorana AM, Samalikova M, Grandori R, Deho G & Polissi A (2011)
990 New insights into the Lpt machinery for lipopolysaccharide transport to the cell surface:
991 LptA-LptC interaction and LptA stability as sensors of a properly assembled
992 transenvelope complex. *J Bacteriol* **193**: 1042-1053.

993 Sriramulu DD, Lunsdorf H, Lam JS & Romling U (2005) Microcolony formation: a novel biofilm
994 model of *Pseudomonas aeruginosa* for the cystic fibrosis lung. *J Med Microbiol* **54**: 667-
995 676.

996 Stead CM, Beasley A, Cotter RJ & Trent MS (2008) Deciphering the unusual acylation pattern
997 of *Helicobacter pylori* lipid A. *J Bacteriol* **190**: 7012-7021.

998 Stead CM, Zhao J, Raetz CR & Trent MS (2010) Removal of the outer Kdo from *Helicobacter*
999 *pylori* lipopolysaccharide and its impact on the bacterial surface. *Mol Microbiol* **78**: 837-
1000 852.

1001 Thaipisuttikul I, Hittle LE, Chandra R, *et al.* (2014) A divergent *Pseudomonas aeruginosa*
1002 palmitoyltransferase essential for cystic fibrosis-specific lipid A. *Mol Microbiol* **91**: 158-
1003 174.

1004 Torres-Cruz J & van der Woude MW (2003) Slipped-strand mispairing can function as a
1005 phase variation mechanism in *Escherichia coli*. *J Bacteriol* **185**: 6990-6994.

1006 Tran AX, Whittimore JD, Wyrick PB, McGrath SC, Cotter RJ & Trent MS (2006) The lipid A 1-
1007 phosphatase of *Helicobacter pylori* is required for resistance to the antimicrobial peptide
1008 polymyxin. *J Bacteriol* **188**: 4531-4541.

1009 Tran AX, Karbarz MJ, Wang X, Raetz CR, McGrath SC, Cotter RJ & Trent MS (2004)
1010 Periplasmic cleavage and modification of the 1-phosphate group of *Helicobacter pylori*
1011 lipid A. *J Biol Chem* **279**: 55780-55791.

1012 Trancassini M, Iebba V, Citerà N, *et al.* (2014) Outbreak of *Achromobacter xylosoxidans* in an
1013 Italian cystic fibrosis center: genome variability, biofilm production, antibiotic resistance,
1014 and motility in isolated strains. *Front Microbiol* **5**: 138.

1015 Traverse CC, Mayo-Smith LM, Poltak SR & Cooper VS (2013) Tangled bank of
1016 experimentally evolved *Burkholderia* biofilms reflects selection during chronic infections.
1017 *Proc Natl Acad Sci U S A* **110**: E250-259.

1018 Trent MS, Pabich W, Raetz CRH & Miller SI (2001) A PhoP/PhoQ-induced lipase (PagL) that
1019 catalyzes 3-O-deacylation of lipid A precursors in membranes of *Salmonella typhimurium*.
1020 *J. Biol. Chem.* **276**: 9083-9092.

1021 Uehlinger S, Schwager S, Bernier SP, Riedel K, Nguyen DT, Sokol PA & Eberl L (2009)
1022 Identification of Specific and Universal Virulence Factors in *Burkholderia cenocepacia*
1023 Strains by Using Multiple Infection Hosts. *Infection and Immunity* **77**: 4102-4110.

- 1024 Vaca-Pacheco S, Paniagua-Contreras GL, Garcia-Gonzalez O & de la Garza M (1999) The
1025 clinically isolated FIZ15 bacteriophage causes lysogenic conversion in *Pseudomonas*
1026 *aeruginosa* PAO1. *Curr Microbiol* **38**: 239-243.
- 1027 Valvano MA (2011) Common Themes in Glycoconjugate Assembly Using the Biogenesis of
1028 O-Antigen Lipopolysaccharide as a Model System. *Biochemistry-Moscow* **76**: 729-735.
- 1029 Valvano MA (2015) Genetics and biosynthesis of lipopolysaccharide. *Molecular Medical*
1030 *Microbiology*, Vol. 1 (Tang Y-W, Sussman M, Liu D, Poxton IR & Schwartzman J,
1031 ed.^eds.), p.^pp. 55-89. Academic Press, Amsterdam.
- 1032 Valvano MA, Messner P & Kosma P (2002) Novel pathways for biosynthesis of nucleotide-
1033 activated glycerol-manno-heptose precursors of bacterial glycoproteins and cell surface
1034 polysaccharides. *Microbiology* **148**: 1979-1989.
- 1035 Valvano MA, Patel KB & Furlong SE (2011) Genetics, biosynthesis and assembly of O
1036 antigen. *Bacterial lipopolysaccharides: Structure, chemical synthesis, biogenesis and*
1037 *interaction with host cells*, (Knirel Y & Valvano MA, ed.^eds.), p.^pp. 275-310. Springer
1038 Verlag Publishing Inc., Wein.
- 1039 van Belkum A, Scherer S, van Leeuwen W, Willemse D, van Alphen L & Verbrugh H (1997)
1040 Variable number of tandem repeats in clinical strains of *Haemophilus influenzae*. *Infect.*
1041 *Immun.* **65**: 5017-5027.
- 1042 van der Woude MW & Bäuml AJ (2004) Phase and antigenic variation in bacteria. *Clin*
1043 *Microbiol Rev* **17**: 581-611, table of contents.
- 1044 Vidigal PG, Dittmer S, Steinmann E, Buer J, Rath P-M & Steinmann J (2014) Adaptation of
1045 *Stenotrophomonas maltophilia* in cystic fibrosis: Molecular diversity, mutation frequency
1046 and antibiotic resistance. *Int J Med Microbiol*.
- 1047 Walsh AG, Matewish MJ, Burrows LL, Monteiro MA, Perry MB & Lam JS (2000)
1048 Lipopolysaccharide core phosphates are required for viability and intrinsic drug resistance
1049 in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **35**: 718-727.
- 1050 Wang G, Ge Z, Rasko DA & Taylor DE (2000) Lewis antigens in *Helicobacter pylori*:
1051 biosynthesis and phase variation. *Mol Microbiol* **36**: 1187-1196.
- 1052 Warren AE, Boulianne-Larsen CM, Chandler CB, et al. (2011) Genotypic and phenotypic
1053 variation in *Pseudomonas aeruginosa* reveals signatures of secondary infection and
1054 mutator activity in certain cystic fibrosis patients with chronic lung infections. *Infect Immun*
1055 **79**: 4802-4818.
- 1056 Watson ME, Burns JL & Smith AL (2004) Hypermutable *Haemophilus influenzae* with
1057 mutations in *mutS* are found in cystic fibrosis sputum. *Microbiology* **150**: 2947-2958.
- 1058 Wattam AR, Foster JT, Mane SP, et al. (2014) Comparative phylogenomics and evolution of
1059 the *Brucellae* reveal a path to virulence. *J Bacteriol* **196**: 920-930.
- 1060 West NP, Sansonetti P, Mounier J, et al. (2005) Optimization of virulence functions through
1061 glucosylation of *Shigella* LPS. *Science* **307**: 1313-1317.
- 1062 Whitfield C & Trent MS (2014) Biosynthesis and export of bacterial lipopolysaccharides. *Annu*
1063 *Rev Biochem* **83**: 99-128.
- 1064 WHO (1994) Schistosomes, liver flukes and *Helicobacter pylori*. IARC Monographs on the
1065 Evaluation of Carcinogenic Risks to Humans. *World Health Organization, International*
1066 *Agency for Research on Cancer* **61**: 1-241.

1067 Wildschutte H, Preheim SP, Hernandez Y & Polz MF (2010) O-antigen diversity and lateral
1068 transfer of the wbe region among *Vibrio splendidus* isolates. *Environ Microbiol* **12**: 2977-
1069 2987.

1070 Yang L, Jelsbak L, Marvig RL, *et al.* (2011) Evolutionary dynamics of bacteria in a human
1071 host environment. *Proc Natl Acad Sci U S A* **108**: 7481-7486.

1072 Zlosnik JE, Mori PY, To D, Leung J, Hird TJ & Speert DP (2014) Swimming motility in a
1073 longitudinal collection of clinical isolates of *Burkholderia cepacia* complex bacteria from
1074 people with cystic fibrosis. *PLoS One* **9**: e106428.

1075 Zlosnik JE, Costa PS, Brant R, *et al.* (2011) Muroid and nonmuroid *Burkholderia cepacia*
1076 complex bacteria in cystic fibrosis infections. *Am J Respir Crit Care Med* **183**: 67-72.
1077
1078
1079
1080

1081 **Table 1. LPS genes altered in *P. aeruginosa* during chronic infections.**

LPS metabolism	Genes	Reference
Lipid A biosynthesis and modification	<i>lpxO2</i>	Cramer <i>et al.</i> , 2011; Yang <i>et al.</i> , 2011
	<i>lpxC</i> , <i>yciK</i>	Cramer <i>et al.</i> , 2011
	<i>pagL</i>	Cigana <i>et al.</i> , 2009; Yang <i>et al.</i> , 2011
Core biosynthesis and modification	<i>rfaD</i> , <i>wapP</i>	Cramer <i>et al.</i> , 2011
Common polysaccharide antigen biosynthesis	<i>wbpZ</i>	Cramer <i>et al.</i> , 2011
O-specific antigen biosynthesis	<i>wbpA</i> , <i>pa5238</i>	Smith <i>et al.</i> , 2006
	<i>wzz</i>	Yang <i>et al.</i> , 2011
	<i>wbpA</i> , <i>wbpB</i> , <i>wbpC</i> , <i>wbpD</i> , <i>wbpE</i> , <i>wbpG</i> , <i>wbpH</i> , <i>wbpI</i> , <i>wbpJ</i> , <i>wbpK</i> , <i>wbpL</i> , <i>wzx</i> , <i>wzy</i> , <i>wzz</i> , <i>pa1385</i>	Warren <i>et al.</i> , 2011
O antigen ligase	<i>waaL</i>	Dettman <i>et al.</i> , 2013

1082

1083

1084

1085 **Table 2. LPS genes altered in *B. dolosa* and *B. cenocepacia* during chronic infections.**1086 The homologous gene in *P. aeruginosa* is also indicated.

1087

Gene or locus	Homologous gene in <i>P. aeruginosa</i> PAO1	Reference
<i>wbaD</i>	-	Lieberman <i>et al.</i> , 2011
YP_834517	<i>rmlB</i>	Traverse <i>et al.</i> , 2013
YP_834518	<i>rmlA</i>	Traverse <i>et al.</i> , 2013
YP_834524	<i>migA</i>	Traverse <i>et al.</i> , 2013
YP_834525	<i>wbpW</i>	Traverse <i>et al.</i> , 2013
YP_834526	<i>gmd</i>	Traverse <i>et al.</i> , 2013
YP_834528	-	Traverse <i>et al.</i> , 2013
YP_834530	<i>wapR</i>	Traverse <i>et al.</i> , 2013
YP_834532	<i>wbpL</i>	Traverse <i>et al.</i> , 2013
YP_834533	<i>wbpM</i>	Traverse <i>et al.</i> , 2013

1088

1089

1090

1091

1092

1093

Legend to Figures

Figure 1. Cell envelope organization of Gram-negative bacteria. The cell envelope of Gram-negative bacteria is characterized by the presence of two lipid bilayers: the outer membrane (OM) and the cytoplasmic membrane (CM), which are separated by the periplasm, containing hydrolytic enzymes, binding proteins, chemoreceptors and the peptidoglycan cell wall. The OM is an asymmetric lipid bilayer. The outer leaflet of the OM contains mainly LPS molecules, which form contacts with integral outer membrane proteins (OMPs). The inner layer of the OM and the lipid layers of the cytoplasmic membrane contain phospholipids and membrane proteins.

Figure 2. Simplified overview of the LPS biosynthesis. Lipid A-Kdo₂ is synthesized on the cytoplasmic surface of the cytoplasmic membrane. The rest of the core is assembled to the lipid A-Kdo₂ and MsbA flips the whole complex to the periplasmic side of the cytoplasmic membrane. The O antigen is synthesized by cytoplasmic membrane-associated enzyme complexes using C55-undecaprenol phosphate (Und-P) as an acceptor for chain assembly and is then flipped to the periplasmic face of the membrane by one of the three pathways: (1) Wzy-dependent, (2) ABC-transporter-dependent, or (3) synthase-dependent. For simplicity, only the ABC-transporter pathway is represented. Once on the periplasmic side, the O antigen is linked to the lipid A-core by the WaaL ligase and the mature LPS molecule is then transported across the periplasm and inserted into the outer leaflet of the outer membrane by the Lpt (LPS transport) system, a complex that spans the Gram-negative cell envelope to deliver LPS to the outer membrane (E). OM, Outer membrane; CM, Cytoplasmic membrane.

Figure 3. Lipid A modifications occurring in *P. aeruginosa* during adaptation to long-term chronic infection. The basic tetra-acylated lipid A structure can be modified by: deacylation by PagL; palmitoylation by PagP; acylation by HtrB; acylation by LpxO; addition of Arap4N by PmrAB on position 1 or 4'; and addition of phosphoethanolamine by ColRS on position 1 or 4'.

1124 **Figure 4. Lewis antigen structures.** *H. pylori* can produce type 1 (based on a β -(1,3)-linked
1125 galactose-GlcNAc sugar backbone) and type 2 (based on a β -(1,4)-linked galactose-GlcNAc
1126 sugar backbone) Lewis antigens. Le^a and Le^x are built by addition of a fucose residue to the
1127 GlcNAc sugar of the type 1 and type 2 backbone, through α -(1,4) or α -(1,3)
1128 linkages, respectively. Le^b and Le^y are built by addition of a fucose residue through α -(1,2)
1129 linkage to Le^a and Le^x structures, respectively. Sialyl-Le^x (SLe^x) is built by addition of a sialyl
1130 group to the Le^x antigen by a α -(2,3) linkage.